

Table 1. Sociodemographic and Clinical Characterization	
Characteristic	n = 73 (%)
Age (years) ± SD	26.5 ± 6.2
Weight (kg) ± SD	67.1 ± 10.7
Height (meters) ± SD	1.58 ± 0.06
BMI (kg/m ²) ± SD	26.9 ± 3.9
Procedure	
- Cesarean section	18 (24.7)
- Cesarean section and Tubal Ligation	15 (20.5)
- Tubal Ligation	40 (54.8)
Previous pregnancies	
- 1-3	60 (82.2)
- ≥4	13 (17.8)
Previous vaginal deliveries	
- 1-3	66 (90.4)
- 4	7 (9.6)
Previous cesarean sections	
- 0	38 (52.1)
- 1-2	29 (39.7)
- ≥3	6 (8.2)
Labor prior to cesarean section	7 (21.2)
Pre-cesarean analgesia	
- Epidural	5 (15.1)
- Intravenous	3 (9.1)
Indication for cesarean section	
- Elective	24 (72.3)
- Emergency	9 (27.7)
Anesthetic technique	
- Spinal	72 (98.6)
- Epidural	1 (1.4)
Neuroaxial opioid	
- Fentanyl (mcg) ± SD	19.2 ± 22.3
- Morphine (mcg) ± SD	54.5 ± 51.4
Surgical time (minutes) ± SD	32.2 ± 18
Complications (Hypotension, nausea, and vomiting)	5 (6.8)
Intraoperative analgesia	
- Diclofenac	50 (68.5)
- Dipyrrone	11 (15.1)
Immediate postoperative analgesia	
- Diclofenac	1 (1.4)
- Dipyrrone	32 (43.8)
Post-anesthesia care unit pain (2 hours) (NRS)	
Mean: 0.37 SD 1.33 Median 0 (IQR 0-0)	
- No pain (0)	65 (89.0)
- Mild (1-3)	5 (6.9)
- Moderate (4-6)	2 (2.7)
- Severe (7-10)	1 (1.4)
Pain at 24 hours (NRS)	
Mean: 4.03 SD 2.73 Median 4 (IQR 3-5)	
- No pain (0)	10 (13.7)
- Mild (1-3)	25 (34.3)
- Moderate (4-6)	23 (31.5)
- Severe (7-10)	15 (20.5)
Pain at 48 hours (NRS)	
Mean: 3.85 SD 2.32 Median 4 (IQR 2-5)	
- No pain (0)	9 (12.3)
- Mild (1-3)	25 (34.3)
- Moderate (4-6)	27 (36.9)
- Severe (7-10)	12 (16.5)

SD: Standard Deviation; BMI: Body Mass Index; NRS: Numeric Rating Scale for pain; IQR: Interquartile Range.